U.S. VISA VACCINATION CHECKLIST Adults (aged 19 and over)

NAME

DATE OF BIRTH

For the attention of the general practitioner: Check the presence of the name and date of birth; update vaccinations and record them in the right-hand column with the date indicated; Sign and stamp each.

		VACCINE NAME, DATE OF VACCINATION, SIGNATURE AND STAMP
Diphtheria, tetanus, pertussis	To get a booster if you have not had one in the past 10 years. If you have ever been vaccinated against pertussis the booster does not need to include it.	
POLIO	Provide proof of 3 doses of Polio. If this is not the case, you should have had a Polio vaccine in the last 12 months. Special case - you need to have had a polio vaccination booster within the last 4-52 weeks if you live or have spent more than 4 weeks in certain countries designated as at high risk. You must consult the list of all current countries affected on the CDC website: https://www.cdc.gov/immigrant-refugee-health/hcp/panel-physicians/polio-addendum.html.	
Hepatitis B	If under 60 years old. 3 doses – the second On D+1 month and third on D+6 months. At least one vaccine before the visit.	
Rubella, mumps, measles (MMR)	To do if you were born from 1957 onwards. At least 1 before the visit. If there is no written proof of vaccination, you should get vaccinated or have a blood test to see if you are immune to the 3 diseases. Ask your doctor for a prescription. If you are only immune to 1 or 2 of these diseases, you must be vaccinated again with MMR. Individual vaccines do not exist. A history of rubella, mumps, measles is not enough as proof. Contraindicated vaccine in pregnancy	
Chickenpox Varicella	If there is no written proof of vaccination, you should get vaccinated or have a blood test to see if you are immune. Ask your doctor for a prescription. A history of chickenpox will only suffice if you have the date noted by a doctor. Contraindicated vaccine in pregnancy	
Seasonal flu	The vaccine is mandatory during the flu season or as long as the vaccine remains available in pharmacies. Check availability with your pharmacist.	
Pneumococcus	The pneumococcal vaccine is mandatory from the age of 65 . U.S. Centers for Disease Control (CDC) calls for first vaccination with PCV20 pneumococcal vaccine	